



## Coordination diversity of *N*-phosphoryl-*N'*-phenylthiourea (LH) towards $\text{Co}^{\text{II}}$ , $\text{Ni}^{\text{II}}$ and $\text{Pd}^{\text{II}}$ cations: Crystal structure of $\text{ML}_2\text{-N,S}$ and $\text{ML}_2\text{-O,S}$ chelates

Felix D. Sokolov<sup>a</sup>, Nail G. Zabirow<sup>a</sup>, Luiza N. Yamaliev<sup>a</sup>, Valery G. Shtyrin<sup>a</sup>,  
Ruslan R. Garipov<sup>a</sup>, Vasily V. Brusko<sup>a</sup>, Alexander Yu. Verat<sup>a</sup>, Sergey V. Baranov<sup>a</sup>,  
Piotr Mlynarz<sup>b,1</sup>, Tadeusz Glowiak<sup>b</sup>, Henryk Kozlowski<sup>b,\*</sup>

<sup>a</sup> Kazan State University, Department of Chemistry, Kremlevskaya Street 18, Kazan 420008, Russia

<sup>b</sup> University of Wrocław, Faculty of Chemistry, F. Joliot-Curie 14, 50-383 Wrocław, Poland

Received 27 September 2005; received in revised form 2 January 2006; accepted 15 January 2006

Available online 28 February 2006

### Abstract

Thiourea,  $\text{PhNHC(S)NHP(O)(OPr}^i)_2$  (LH) chelates of  $\text{Co}^{\text{II}}$ ,  $\text{Ni}^{\text{II}}$ , and  $\text{Pd}^{\text{II}}$  ions have been obtained and investigated by single-crystal X-ray diffraction, UV, IR, NMR spectroscopy, and EI mass-spectrometry. The unusual 1,3-*N,S*-coordination via sulfur and NP(O) nitrogen atoms has been found in the *trans*-square-planar  $\text{NiL}_2$  and  $\text{PdL}_2$  complexes, whereas the 1,5-*O,S*-coordination is realized in the tetrahedral  $\text{CoL}_2$  complex. DFT calculations have revealed significant stabilization of the 1,3-*N,S*-structures due to stronger crystal field and the  $\text{NH-O=P}$  hydrogen bonds.

© 2006 Elsevier B.V. All rights reserved.

**Keywords:** Thiourea coordination; Substituent effects;  $\text{Co}^{\text{II}}$  complexes;  $\text{Ni}^{\text{II}}$  complexes;  $\text{Pd}^{\text{II}}$  complexes; *N,S*-coordination; *O,S*-coordination

### 1. Introduction

$\beta$ -Dicarbonyl compounds (*N*-acylureas **1**, acetylacetonates, etc.) and their phosphorus containing analogs (*N*-acylamidophosphinates **2**, imidodiphosphinates **3**) (Fig. 1) were found to be very attractive ligands for variety of metal ions [1–4].

$\beta$ -Bifunctional ligands **1–3** contain three potential donor centers: X, Y and internal nitrogen atoms. Nucleophilic ability of the latter donor is significantly reduced by the two neighboring electron-withdrawing groups. That is why the alkylation of *N*-thioacylamido(thio)phosphates-

$\text{RC(S)NHP(Y)(OR}^i)_2$  ( $\text{Y} = \text{O,S}$ ) always proceeds on the thiocarbonyl sulfur atom [1a,5].

There is a fair amount of the data concerning the structures of complexes of divalent transition metal ions of Ib, IIb, VIIIb groups with *N*-thioacylamidothiophosphinates (**2**) ( $\text{X, Y} = \text{S}$ ) [1–5,7–10]. The 1,5-bidentate coordination of deprotonated ligands through sulfur atoms takes place in all these cases.

The substituents (or other factors) are hardly capable of changing a coordination mode in the complexes of *dithio*-derivatives of **2** with divalent cations of VIIIb group. Due to the strong  $\pi$ -interactions, the formation of metal–sulfur bonds is most energetically favorable in all these cases.

Replacement of one sulfur atom in conjugated  $[\text{RC(X)NP(Y)R}_2']^{(-)}$  moiety to oxygen ( $\text{X} = \text{S, Y} = \text{O}$  or  $\text{X} = \text{O, Y} = \text{S}$ ) leads to appearance of the ambivalent coordination modes and possible competition between the three donor centers in the binding anion species. As a result, the

\* Corresponding author.

E-mail addresses: [felix.sokolov@ksu.ru](mailto:felix.sokolov@ksu.ru) (F.D. Sokolov), [henrykoz@wchuwr.chem.uni.wroc.pl](mailto:henrykoz@wchuwr.chem.uni.wroc.pl) (H. Kozlowski).

<sup>1</sup> Present address: Wrocław University of Technology, Wybrzeże Wyspiańskiego 27, 50-370 Wrocław, Poland.